

23



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : A61K 31/35, 31/28, 31/05, 31/34, 31/44, 31/40, 31/195</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/42332 (43) International Publication Date: 1 October 1998 (01.10.98)</p>
<p>(21) International Application Number: PCT/US98/05414 (22) International Filing Date: 19 March 1998 (19.03.98) (30) Priority Data: 60/040,673 21 March 1997 (21.03.97) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indi- anapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FLEISCH, Jerome, H. [US/US]; 10532 Coppergate, Carmel, IN 46032 (US). JACKSON, William, T. [US/US]; 7036 Bexley Drive, Indianapolis, IN 46256 (US). SAWYER, Jason, S. [US/US]; 5718 North Winthrop Avenue, Indianapolis, IN 46220 (US). (74) Agents: LENTZ, Nelsen, L. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	
<p>(54) Title: LEUKOTRIENE ANTAGONISTS USEFUL FOR TREATING IRITIS</p> <p>(57) Abstract</p> <p>This invention provides methods for the treatment or inhibiting of iritis which comprises administering to a mammal in need thereof an effective amount of a compound having activity as a leukotrienne B<sub>4</sub> antagonist.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

## LEUKOTRIENE ANTAGONISTS USEFUL FOR TREATING IRITIS

A red eye is a common complaint, often related to  
5 benign conditions. However, a red eye in conjunction with  
symptoms such as photophobia, pain and decreased visual  
acuity may be a much more serious disorder.

Intraocular inflammation may result in deleterious  
structural alterations such as cataracts, synchiae and  
10 glaucoma. It may also lead to microvascular leakage in the  
retina circulation.

Iritis or inflammation of the iris and conjunctivitis  
can be precipitated by a variety of causes. For example, in  
addition to infectious causes, allergies and hypersensitivity  
15 may cause iritis and it occurs as a secondary complication  
in five to ten percent of patients with ulcerative colitis,  
and up to 30% of patients with ankylosing spondylitis.

Once the diagnosis is made, treatment with mydriatic  
and cycloplegic agents with topical corticosteroids is  
20 imperative. However, close monitoring is required because  
overuse of corticosteroids has ominous side effects.

Research in the area of allergic reactions of the lung  
has provided evidence that arachidonic acid derivatives  
formed by the action of lipxygenases are related to various  
25 disease states. Some of these arachidonic acid metabolites  
have been classified as members of a family of  
eicosatetraenoic acids termed leukotrienes. Three of these  
substances are currently thought to be major components of  
what has been previously called slow reacting substance of  
30 anaphylaxis (SRS-A) and have been designated leukotrienes  
C<sub>4</sub>, D<sub>4</sub>, and E<sub>4</sub> (LTC<sub>4</sub>, LTD<sub>4</sub>, and LTE<sub>4</sub>, respectively).


Another arachidonic acid metabolite, leukotriene B<sub>4</sub>  
(LTB<sub>4</sub>), is a proinflammatory lipid which has been implicated  
in the pathogenesis of psoriasis, arthritis, chronic lung  
35 diseases, acute respiratory distress syndrome, shock,  
asthma, inflammatory bowel diseases, and other inflammatory  
states characterized by the infiltration and activation of

-2-

polymorphonuclear leukocytes and other proinflammatory cells. Thus, when activated, the polymorphonuclear leukocytes liberate tissue-degrading enzymes and reactive chemicals causing the inflammation. Antagonism of LTB<sub>4</sub> should therefore provide a novel therapeutic approach to treatment of these and other LTB<sub>4</sub> mediated conditions.

Because of the debilitating effects of ophthalmic disorders such as iritis there continues to exist a need for effective treatments.

This invention provides a method for the treatment or inhibiting of iritis in mammals comprising administering to a mammal in need thereof an effective amount of a compound of Formula I



wherein:

R<sub>1</sub> is C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>2</sub>-C<sub>5</sub> alkynyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, (C<sub>1</sub>-C<sub>4</sub> alkyl)thio, halo, or R<sub>2</sub>-substituted phenyl;


each R<sub>2</sub> and R<sub>3</sub> are each independently hydrogen, halo, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(O)<sub>q</sub>-, trifluoromethyl, or di-(C<sub>1</sub>-C<sub>3</sub> alkyl)amino;

X is -O-, -S-, -C(=O), or -CH<sub>2</sub>-;

Y is -O- or -CH<sub>2</sub>-;

-3-

or when taken together, -X-Y- is -CH=CH- or





5

Z is a straight or branched chain C<sub>1</sub>-C<sub>10</sub> alkyldidenyl;


10

A is a bond, -O-, -S-, -CH=CH-, or -CR<sub>a</sub>R<sub>b</sub>-, where R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, C<sub>1</sub>-C<sub>5</sub> alkyl, or R<sub>7</sub>-substituted phenyl, or when taken together with the carbon atom to which they are attached form a C<sub>4</sub>-C<sub>8</sub> cycloalkyl ring;




R<sub>4</sub> is R<sub>6</sub>



15



-4-



where,

5

each R<sub>6</sub> is independently -COOH, 5-tetrazolyl, -CON(R<sub>9</sub>)<sub>2</sub>, or -CONHSO<sub>2</sub>R<sub>10</sub>;

10

each R<sub>7</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>2</sub>-C<sub>5</sub> alkynyl, benzyl, methoxy, -W-R<sub>6</sub>, -T-G-R<sub>6</sub>, (C<sub>1</sub>-C<sub>4</sub> alkyl)-T-(C<sub>1</sub>-C<sub>4</sub> alkylidenyl)-O-, or hydroxy;

R<sub>8</sub> is hydrogen or halo;

-5-

each R<sub>9</sub> is independently hydrogen, phenyl, or C<sub>1</sub>-C<sub>4</sub> alkyl, or when taken together with the nitrogen atom form a morpholino, piperidino, piperazino, or pyrrolidino group;

5

R<sub>10</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl or phenyl;

R<sub>11</sub> is R<sub>2</sub>, -W-R<sub>6</sub>, or -T-G-R<sub>6</sub>;

10

each W is a bond or straight or branched chain divalent hydrocarbyl radical of one to eight carbon atoms;

15

each G is a straight or branched chain divalent hydrocarbyl radical of one to eight carbon atoms;

each T is a bond, -CH<sub>2</sub>-, -O-, -NH-, -NHCO-, -C(=O)-, or -S(O)<sub>q</sub>-;

20

K is -C(=O)- or -CH(OH)-;

each q is independently 0, 1, or 2;

p is 0 or 1; and

25

t is 0 or 1;

provided when X is -O- or -S-, Y is not -O-;

30

provided when A is -O- or -S-, R<sub>4</sub> is not R<sub>6</sub>;

provided when A is -O- or -S- and Z is a bond, Y is not -O-; and

35

provided W is not a bond when p is 0;

-6-

or a pharmaceutically acceptable salt or solvate thereof.

The following definitions refer to the various terms used throughout this disclosure.

The term "C<sub>1</sub>-C<sub>5</sub> alkyl" refers to the straight and  
5 branched aliphatic radicals of 1 to 5 carbon atoms such as methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, n-pentyl, 2,2-dimethylpropyl, and the like. Included within this definition are the terms "C<sub>1</sub>-C<sub>3</sub> alkyl" and "C<sub>1</sub>-C<sub>4</sub> alkyl".

10 The term "C<sub>2</sub>-C<sub>5</sub> alkenyl" refers to straight and branched aliphatic radicals of 2 to 5 carbon atoms containing one double bond, such as -CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>, and the like.

The term "C<sub>2</sub>-C<sub>5</sub> alkynyl" refers to straight and  
15 branched aliphatic residues of 2 to 5 carbon atoms containing one triple bond, such as -C≡CH, -CH<sub>2</sub>-C≡CH, -CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH(CH<sub>3</sub>)C≡CH, -CH<sub>2</sub>C≡CCH<sub>3</sub>, and the like.

The term "C<sub>1</sub>-C<sub>4</sub> alkoxy" refers to methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, and tert-butoxy.

20 The term "halo" refers to fluoro, chloro, bromo, and iodo.

The term "C<sub>1</sub>-C<sub>10</sub> alkylidenyl" refers to a divalent radical derived from a C<sub>1</sub>-C<sub>10</sub> alkane such as -CH<sub>2</sub>-,  
-CH(CH<sub>3</sub>)-, -C(CH<sub>3</sub>)<sub>2</sub>-, -CH(C<sub>2</sub>H<sub>5</sub>)-, -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH(CH<sub>3</sub>)-,  
25 -CH(CH<sub>3</sub>)CH<sub>2</sub>-, -CH(CH<sub>3</sub>)CH(CH<sub>3</sub>)-, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>-, -CH<sub>2</sub>CH(C<sub>2</sub>H<sub>5</sub>)-,  
-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH(CH<sub>3</sub>)CH<sub>2</sub>-,  
-CH<sub>2</sub>CH(C<sub>2</sub>H<sub>5</sub>)CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH(C<sub>2</sub>H<sub>5</sub>)-, -C(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-,  
-CH(CH<sub>3</sub>)CH<sub>2</sub>CH(CH<sub>3</sub>)-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-,  
-CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH(C<sub>2</sub>H<sub>5</sub>)CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-,  
30 -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>10</sub>-, and the like. Included within this definition are the terms "C<sub>1</sub>-C<sub>4</sub> alkylidene" and "C<sub>2</sub>-C<sub>4</sub> alkylidene".

The term "C<sub>4</sub>-C<sub>8</sub> cycloalkyl" refers to a cycloalkyl  
ring of four to eight carbon atoms, such as cyclobutyl,  
35 cyclopentyl, cyclohexyl, 4,4-dimethylcyclohexyl,  
cycloheptyl, cyclooctyl, and the like.



-7-

The term "straight or branched chain divalent hydrocarbyl residue of one to eight carbon atoms" refers to a divalent radical derived from a straight or branched alkane, alkene, or alkyne of one to eight carbon atoms.

5 Depending upon the branching and number of carbon atoms, as will be appreciated by organic chemists, such a moiety can contain one, two or three double or triple bonds, or combinations of both. As such, this term can be considered an alkylidene group as defined above containing from 1 to 8  
10 carbon atoms optionally containing one to three double or triple bonds, or combinations of the two, limited as noted in the preceding sentence.

This invention includes the pharmaceutically acceptable base addition salts of the compounds of Formula  
15 I. Such salts include those derived from inorganic bases, such as ammonium and alkali and alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, as well as salts derived from basic organic amines, such as aliphatic and aromatic amines, aliphatic diamines, hydroxy  
20 alkylamines, and the like. Such bases useful in preparing the salts of this invention thus include ammonium hydroxide, potassium carbonate, sodium bicarbonate, calcium hydroxide, methyl amine, diethyl amine, ethylene diamine, cyclohexylamine, ethanolamine, and the like. The potassium  
25 and sodium salt forms are particularly preferred.


This invention includes both mono-salt forms, i.e., a 1:1 ratio of a compound of Formula I with a base as previously described, as well as di-salt forms in those instances where a compound of Formula I has two acidic  
30 groups. In addition, this invention includes any solvate forms of the compounds of Formula I or salts thereof, such as ethanol solvates, hydrates, and the like.

It is recognized that in compounds having branched alkyl, alkylidenyl, or hydrocarbyl functionality, and in  
35 those compounds bearing double or triple bonds, various stereoisomeric products may exist. This invention is not limited to any particular stereoisomer but includes all

-8-

possible individual isomers and mixtures thereof. The term "5-tetrazolyl" refers to both tautomers, ie, (1H)-5-tetrazolyl and (2H)-5-tetrazolyl.

A most preferred group of compounds employed in the methods of the present invention are those compounds of Formula Ia:



10

and pharmaceutically acceptable base addition salts thereof. Especially preferred are those compounds wherein R<sub>2</sub> is halo, particularly fluoro. Preferred R<sub>1</sub> substituents are propyl and especially ethyl.

15

Preferred Z substituents include C<sub>2</sub>-C<sub>4</sub> alkylidene, particularly -CH<sub>2</sub>CH<sub>2</sub>- and -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-. Preferred A groups include -O-, -CH<sub>2</sub>-, -CH(R<sub>7</sub>-substituted phenyl)-, and -C(CH<sub>3</sub>)<sub>2</sub>-.

20

Preferred R<sub>4</sub> groups include -COOH, 5-tetrazolyl, or a mono-, di-, or tri-cyclic group as drawn above wherein there is at least one acidic group attached to a ring, such as -W-COOH, -T-G-COOH, or the corresponding tetrazole derivatives. The preferred W moiety is that of a bond or straight chain C<sub>1</sub>-C<sub>4</sub> alkylidene; preferred G moieties are straight chain C<sub>1</sub>-C<sub>4</sub> alkylidene. It is preferred that R<sub>5</sub> or R<sub>7</sub> be C<sub>1</sub>-C<sub>4</sub> alkyl, especially n-propyl.


25

Particularly preferred groups are those wherein A is -CH(R<sub>7</sub>-substituted phenyl)- and R<sub>4</sub> is -COOH or 5-tetrazolyl. Also preferred are those compounds wherein A is

30

-O- and R<sub>4</sub> is

-9-



Preferred aspects of this substructure are those wherein R<sub>7</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl, especially n-propyl, and R<sub>6</sub> is  
 5 -W-COOH. Particularly preferred are those compounds wherein T is -O- or -S- and W is a bond.

Particularly preferred compounds of the instant invention include 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid;  
 10 3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)-6-(4-carboxyphenoxy)phenyl)propionic acid; 1-(4-(carboxymethoxy)phenyl)-1-(1H-tetrazol-5-yl)-6-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)hexane ; 3-[4-[7-carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]-9H-xanthene]]propanoic acid; 5-[3-[2-(1-carboxy)-ethyl]-4-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenyl]-4-pentynoic acid or a pharmaceutically  
 15 acceptable salt or solvate thereof.

20 The leukotriene B<sub>4</sub> (LTB<sub>4</sub>) antagonists employed in the methods of the present invention may be synthesized essentially as described in US Patent No. 5,462,954 issued October 31, 1995, the entire contents of which are herein incorporated by reference.

25 The following examples further illustrate the preparation of the intermediates and compounds employed in this invention. The examples are illustrative only and are not intended to limit the scope of the invention. Melting points were determined on a Thomas-Hoover apparatus and are uncorrected. NMR spectra were determined on a GE QE-300  
 30 spectrometer. All chemical shifts are reported in parts per million (ppm) relative to tetramethylsilane. Chemical shifts


-10-

of aromatic protons of quinoline species in DMSO-d<sub>6</sub> are concentration dependent. The following abbreviations are used to denote signal patterns: s = singlet, d = doublet, t = triplet, q = quartet, b = broad, m = multiplet. Infrared spectra were determined on a Nicolet DX10 FT-IR spectrometer. Mass spectral data were determined on a CEC-21-110 spectrometer using electron impact (EI) conditions, a MAT-731 spectrometer using free desorption (FD) conditions, or a VG ZAB-3F spectrometer using fast atom bombardment (FAB) conditions. Silica gel chromatography was performed using ethyl acetate/hexane gradients unless otherwise indicated. Reverse-phase chromatography was performed on MCI CHP20P gel using an acetonitrile/water or methanol/water gradient unless otherwise indicated. Tetrahydrofuran (THF) was distilled from sodium/benzophenone ketyl immediately prior to use. All reactions were conducted under argon atmosphere with stirring unless otherwise noted. Where structures were confirmed by infra-red, proton nuclear magnetic resonance, or mass spectral analysis, the compound is so designated by "IR", "NMR", or "MS", respectively.

Example 1

3-[2-[3-[(5-Ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-1-dibenzofuran]propanoic acid disodium salt

-11-



A. Preparation of 3,3-diethoxy-2,3-dihydro-1H-benzofuro-[3,2-f][1]benzopyran.

5

A solution of 2-hydroxydibenzofuran (5.00 g, 27.2 mmol), triethylorthoacrylate (10.1 g, 54.3 mmol) and pivalic acid (1.39 g, 13.6 mmol) in toluene (100 mL) was refluxed for 18 hours. The mixture was cooled to room temperature and washed once with water and once with a saturated sodium bicarbonate solution, dried over sodium sulfate, filtered and concentrated *in vacuo* to provide an orange oil. This material was diluted with hexane and maintained at -20°C for 18 hours. The resulting crystals were collected via vacuum filtration to provide 5.67 g (67%) of the desired title intermediate, mp 64°C; NMR (CDCl<sub>3</sub>) 7.96 (d, J = 7.8 Hz, 1H), 7.57 (d, J = 8.0 Hz, 1H), 7.46 (t, J = 8 Hz, 1H), 7.35 (m, 2H), 7.06 (d, J = 8.8 Hz, 1H), 3.82 (q, J = 7.2 Hz, 2H), 3.73 (q, J = 6.8 Hz, 2H), 3.35 (t, J = 6.9 Hz, 2H), 2.29 (t, J = 7.0 Hz, 2H), 1.23 (t, J = 7.1 Hz, 6H); MS-FD m/e 312

10

15

20

-12-

(p); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ) 2982, 1494, 1476, 1451, 1434, 1251, 1090, 1054, 975.

Analysis for  $\text{C}_{19}\text{H}_{20}\text{O}_4$ :

Calc: C, 73.06; H, 6.45;  
5 Found: C, 72.81; H, 6.72.

B. Preparation of 3-[1-(2-hydroxydibenzofuran)]-propanoic acid ethyl ester.

10 A mixture of 3,3-diethoxy-2,3-dihydro-1H-benzofuro-[3,2-f][1]benzopyran (3.50 g, 11.2 mmol) and 10% aqueous hydrochloric acid (5 mL) in ethyl acetate (30 mL) was stirred at room temperature for 1 hour. The resulting mixture was washed once with water, dried over sodium sulfate, filtered  
15 and concentrated in vacuo to provide a tan solid. Recrystallization from hexane/ethyl acetate provided 3.11 g (98%) of the desired title intermediate as an off-white crystalline material: mp 128-131°C; NMR ( $\text{CDCl}_3$ ) 7.88 (d, J = 7.7 Hz, 1H), 7.59 (d, J = 8.4 Hz, 1H), 7.47 (t, J = 7.2 Hz, 1H), 7.37 (d, J = 8.9 Hz, 1H), 7.36 (t, J = 6.6 Hz, 1H), 7.13 (d, J = 8.8 Hz, 1H), 7.13 (q, J = 8.8 Hz, 2H), 3.43 (t, J = 5.8 Hz, 2H), 3.01 (t, J = 7.7 Hz, 2H), 1.23 (t, J = 7.2 Hz, 3H); MS-FD m/e 284 (100, p), 256 (65), 238 (17); IR (KBr,  $\text{cm}^{-1}$ ) 2985 (b), 1701, 1430, 1226, 1183, 1080.

25 Analysis for  $\text{C}_{17}\text{H}_{16}\text{O}_4$ :

Calc: C, 71.82; H, 5.67;  
Found: C, 71.90; H, 5.43.

C. Preparation of 3-[2-[3-[[5-ethyl-2-(phenylmethoxy)-[1,1'-biphenyl]-4-yl]oxy]propoxy]-1-dibenzofuran]propanoic acid ethyl ester.

3-[1-(2-Hydroxydibenzofuran)]propanoic acid ethyl ester (625 mg, 2.20 mmol) was dissolved in dimethylformamide (10 mL) and carefully treated at room temperature with 95% sodium hydride (58 mg, 2.4 mmol). When gas evolution had

-13-

ceased, 2-benzyloxy-1-phenyl-5-ethyl-4-(3-chloro-1-propyloxy)benzene (836 mg, 2.20 mmol) was added and the resulting mixture was stirred for 18 hours. The mixture was diluted with ether and washed once with water. The organic layer was dried over sodium sulfate, filtered, and concentrated in vacuo to provide a dark oil. Silica gel chromatography (ethyl acetate/hexane) provided 200 mg (14%) of the desired titled intermediate as a colorless oil: NMR (CDCl<sub>3</sub>) 8.11 (d, J = 7.7 Hz, 1H), 7.57 (m, 3H), 7.48 (t, J = 7.3 Hz, 1H), 7.20-7.44 (m, 10 H), 7.17 (s, 1H), 7.08 (d, J = 8.9 Hz, 1H), 6.67 (s, 1H), 5.05 (s, 2H), 4.29 (t, J = 6.2 Hz, 2H), 4.26 (t, J = 6.1 Hz, 2H), 4.15 (q, J = 7.2 Hz, 2H), 3.54 (t, J = 8.5 Hz, 2H), 2.67 (m, 4H), 2.37 (t, J = 6.0 Hz, 2H), 1.21 (m, 6H).

D. Preparation of 3-[2-[3-[(5-ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-1-dibenzofuran]propanoic acid disodium salt.

To a nitrogen-purged solution of 3-[2-[3-[(5-ethyl-2-(phenylmethoxy)[1,1'-biphenyl]-4-yl)oxy]propoxy]-1-dibenzofuran]propanoic acid ethyl ester (200 mg, 0.318 mmol) in a 1:1 mixture of methanol/tetrahydrofuran (40 mL) was added 10% palladium on carbon (25 mg). The resulting suspension was hydrogenated at 1 atm pressure for 24 hours at room temperature. The mixture was filtered through a short pad of Florisil® and the filtrate concentrated in vacuo. The residue was dissolved in a 1:1 mixture of methanol/tetrahydrofuran (20 mL) and treated with 5N sodium hydroxide solution (2 mL) at room temperature for 24 hours. The resulting mixture was extracted once with diethyl ether. The aqueous layer was acidified with 5N hydrochloric acid solution and extracted twice with methylene chloride. The combined methylene chloride fractions were concentrated in vacuo. The residue was dissolved in a minimum of 1N sodium hydroxide solution and purified on HP-20 resin to provide 53 mg (30%) of the desired title product as a fluffy white

-14-

solid: NMR (DMSO- $d_6$ ) 8.12 (d,  $J$  = 6.9 Hz, 1H), 7.64 (d,  $J$  = 8.2 Hz, 1H), 7.37-7.57 (m, 5H), 7.30 (m, 2H), 7.14 (m, 2H), 6.96 (s, 1H), 6.93 (s, 1H), 4.30 (t,  $J$  = 7.3 Hz, 2H), 4.14 (t,  $J$  = 5.4 Hz, 2H), 2.48 (m, 4H), 2.23 (m, 4H), 1.10 (t,  $J$  = 7.6 Hz, 3H); MS-FAB  $m/e$  555 (88,  $p + 1$ ), 533 (62); IR (CHCl<sub>3</sub>,  $cm^{-1}$ ) 3384 (b), 2969, 1566, 1428, 1257, 1181.

Analysis for C<sub>32</sub>H<sub>28</sub>O<sub>6</sub>Na<sub>2</sub>:

Calc: C, 69.31; H, 5.09;


Found: C, 69.51; H, 5.39.

10

### Example 2

7-Carboxy-9-oxo-3-[3-(2-ethyl-5-hydroxy-4-phenylphenoxy)propoxy]-9H-xanthene-4-propanoic acid disodium salt monohydrate

15



A mixture of 2-benzyloxy-1-phenyl-5-ethyl-4-(3-chloro-1-propyloxy)benzene (749 mg, 1.97 mmol), ethyl 7-carboethoxy-3-hydroxy-9-oxo-9H-xanthene-4-propanoate (729 mg, 1.97 mmol), potassium carbonate (1.36 g, 9.85 mmol) and potassium iodide (33 mg, 0.20 mmol) was refluxed for 24 hours. Dimethylsulfoxide (2 mL) was added and heating continued for 24 hours. The reaction mixture was cooled to room temperature, diluted with ethyl acetate, and washed once with water. The organic layer was dried over sodium sulfate, filtered and concentrated in vacuo to reveal a tan solid. This material was dissolved in ethyl acetate (30 mL) and the resulting solution purged with nitrogen. To this solution was added 10% palladium on carbon (120 mg) and the

30



-15-

resulting suspension hydrogenated at 1 atmosphere of pressure. The solution was filtered and concentrated in vacuo to provide a colorless oil. This material was dissolved in a solution of 1:1 methanol/tetrahydrofuran (30 mL) and treated with 5N sodium hydroxide solution (2 mL) at room temperature for 18 hours. The resulting solution was extracted once with diethyl ether and the aqueous layer acidified with 5N hydrochloric acid solution. The resulting precipitate was collected via suction filtration. This material was converted to the di-sodium salt and purified as described above for the preparation of Example 1(D) to provide 390 mg (56%) of the desired title product as a fluffy white solid: NMR (DMSO- $d_6$ ) 12.65 (s, 1H, -OH), 8.65 (s, 1H), 8.28 (dd, J = 8.5, 2.0 Hz, 1H), 8.01 (d, J = 8.9 Hz, 1H), 7.50 (m, 3H), 7.29 (t, J = 7.8 Hz, 2H), 7.17 (m, 2H), 6.93 (s, 1H), 6.89 (s, 1H), 4.26 (m, 4H), 3.12 (m, 2H), 2.47 (m, 2H), 2.23 (m, 2H), 1.10 (t, J = 7.4 Hz, 3H); MS-FAB m/e 627 (24, p), 605 (40), 583 (24), 331 (24), 309 (100); IR (KBr,  $cm^{-1}$ ) 3419 (b), 2962, 1612, 1558, 1443, 1390, 1277, 1084.

Analysis for  $C_{34}H_{28}O_9Na_2 \cdot H_2O$ :

Calc: C, 63.34; H, 4.69;


Found: C, 63.36; H, 4.50.

25

Example 3

2-[2-Propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid sodium salt

-16-



A. Preparation of 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-(phenylmethoxy)phenoxy]propoxy]phenoxy]-5-benzoic acid methyl ester.

A mixture of 2-benzyloxy-1-(4-fluorophenyl)-5-ethyl-4-(3-chloro-1-propyloxy)benzene (20.0 g, 50.2 mmol) and sodium iodide (75.3 g, 502 mmol) in 2-butanone (200 mL) was refluxed for 6 hours. The mixture was diluted with ether and washed once with water. The organic layer was dried over sodium sulfate, filtered, and concentrated in vacuo to provide a colorless oil. This material was dissolved in

-17-

dimethylformamide (100 mL) and treated with 2-(3-hydroxy-2-propylphenoxy)benzoic acid methyl ester (14.4 g, 50.2 mmol) and potassium carbonate (20.8 g, 151 mmol) at room temperature for 24 hours. This mixture was diluted with  
5 water and twice extracted with ether. The aqueous layer was separated and back-extracted once with ethyl acetate. The combined organic layers were dried over sodium sulfate, filtered, and concentrated in vacuo to provide a yellow oil. Silica gel chromatography provided 25.4 g (78%) of the  
10 desired title intermediate as a pale golden oil: NMR (CDCl<sub>3</sub>) 7.91 (d, J = 7.8 Hz, 1H), 7.54 (d, J = 8.6 Hz, 1H), 7.52 (d, J = 8.5 Hz, 1H), 7.25-7.43 (m, 6H), 7.03-7.38 (m, 5H), 6.84 (d, J = 8.3 Hz, 1H), 6.71 (d, J = 8.1 Hz, 1H), 6.63 (s, 1H), 6.47 (d, J = 8.1 Hz, 1H), 5.03 (s, 2H), 4.24  
15 (t, J = 5.7 Hz, 2H), 4.21 (t, J = 5.8 Hz, 2H), 3.86 (s, 3H), 2.69 (t, J = 7.8 Hz, 2H), 2.64 (t, J = 7.7 Hz, 2H), 2.34 (quintet, J = 6.0 Hz, 2H), 1.60 (hextet, J = 5.0 Hz, 2H), 1.22 (t, J = 7.5 Hz, 3H), 0.94 (t, J = 7.5 Hz, 3H); MS-FD m/e 648 (p); IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 2960, 1740, 1604, 1497, 1461,  
20 1112.

Analysis for C<sub>41</sub>H<sub>41</sub>O<sub>6</sub>F:

Calc: C, 75.91; H, 6.37;

Found: C, 76.15; H, 6.45.

25 B. Preparation of 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid methyl ester.

2-[2-Propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-(phenylmethoxy)phenoxy]propoxy]phenoxy]benzoic acid methyl  
30 ester (33.0 g, 50.9 mmol) was de-benzylated as described above for the preparation of Example 2 to provide 27.3 g (96%) of the title intermediate as an amber oil: NMR (CDCl<sub>3</sub>) 7.90 (dd, J = 7.8, 1.7 Hz, 1H), 7.42 (m, 3H), 7.05-  
35 7.23 (m, 4H), 6.99 (s, 1H), 6.84 (d, J = 8.1 Hz, 1H), 6.70 (d, J = 8.1 Hz, 1H), 6.55 (s, 1H), 6.46 (d, J = 8.1 Hz, 1H),

-18-

5.05 (s, 1H, -OH), 4.23 (m, 4H), 3.86 (s, 3H), 2.68 (t, J = 7.4 Hz, 2H), 2.62 (q, J = 7.5 Hz, 2H), 2.36 (quintet, J = 6.0 Hz, 2H), 1.60 (hextet, J = 7.7 Hz, 2H), 1.20 (t, J = 7.6 Hz, 3H), 0.94 (t, J = 7.4 Hz, 3H); MS-FD m/e 558 (p); IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 2965, 1727, 1603, 1496, 1458, 1306, 1112.

Analysis for C<sub>34</sub>H<sub>35</sub>O<sub>6</sub>F:

Calc: C, 73.10; H, 6.31;

Found: C, 73.17; H, 6.42.

10 C. Preparation of 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid sodium salt.

2-[2-Propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid methyl ester (21.5 g, 38.5 mmol) was hydrolyzed as described above for the preparation of Example 2. The acid was converted to the sodium salt and purified as described above for the preparation of Example 1(D) to provide 16.7 g (77%) of the desired title product as a white amorphous solid: NMR (DMSO-d<sub>6</sub>) 10.50 (bs, 1H, -OH), 7.51 (m, 3H), 7.20 (t, J = 7.4 Hz, 1H), 7.13 (m, 2H), 7.00 (m, 2H), 6.95 (s, 1H), 6.67 (dd, J = 8.2, 3.3 Hz, 2H), 6.62 (s, 1H), 6.26 (d, J = 8.2 Hz, 1H), 4.14 (t, J = 5.8 Hz, 2H), 4.02 (t, J = 5.7 Hz, 2H), 2.60 (t, J = 6.8 Hz, 2H), 2.47 (q, J = 7.3 Hz, 2H), 2.16 (t, J = 5.9 Hz, 2H), 1.45 (hextet, J = 7.5 Hz, 2H), 1.07 (t, J = 7.5 Hz, 3H), 0.81 (t, J = 7.4 Hz, 3H); MS-FAB m/e 568 (38, p + 1), 567 (100, p), 544 (86), 527 (77), 295 (65), 253 (45); IR (KBr, cm<sup>-1</sup>) 3407 (b), 2962, 1603, 1502, 1446, 1395, 1239, 1112.

Analysis for C<sub>33</sub>H<sub>32</sub>O<sub>6</sub>FNa:

Calc: C, 69.95; H, 5.69; F, 3.35;

Found: C, 69.97; H, 5.99; F, 3.52.

35 The methods of the present invention describe the use of leukotriene antagonists for the treatment or

-19-

inhibition of iritis which is characterized by the excessive release of leukotriene B<sub>4</sub>.

The term "excessive release" of a leukotriene refers to an amount of the leukotriene sufficient to cause  
5 iritis. The amount of leukotriene which is considered to be excessive will depend on a variety of factors, including the amount of leukotriene required to cause the disease, and the species of the mammal involved. As will be appreciated by those skilled in the art, the success of treating a mammal  
10 suffering from or susceptible to iritis characterized by an excessive release of leukotriene with a compound of Formula I will be measured by the regression or prevention of the symptoms of the condition.

15

#### Assays

##### Assay 1

The effectiveness of compounds of Formula I to inhibit the binding of tritiated LTB<sub>4</sub> to guinea pig lung membranes was determined as follows.

20

##### [<sup>3</sup>H]-LTB<sub>4</sub> Radioligand Binding Assay in Guinea Pig Lung Membranes

[<sup>3</sup>H]-LTB<sub>4</sub> (196-200 Ci/mmol) was purchased from New  
25 England Nuclear (Boston, MA). All other materials were purchased from Sigma (St. Louis, MO). Incubations (555 mL) were performed in polypropylene minitubes for 45 minutes at 30°C and contained 25 mg of guinea pig lung membrane protein (Silbaugh, et al., European Journal of Pharmacology, 223  
30 (1992) 57-64) in a buffer containing 25 mM MOPS, 10 mM MgCl<sub>2</sub>, 10 mM CaCl<sub>2</sub>, pH 6.5, approximately 140 pM [<sup>3</sup>H]-LTB<sub>4</sub>, and displacing ligand or vehicle (0.1% DMSO in 1 mM sodium carbonate, final concentration) as appropriate. The binding reaction was terminated by the addition of 1 mL ice cold  
35 wash buffer (25 mM Tris-HCl, pH 7.5) followed immediately by vacuum filtration over Whatman GF/C glass fiber filters using a Brandel (Gaithersburg, MD) 48 place harvester. The

-20-

filters were washed three times with 1 mL of wash buffer. Retained radioactivity was determined by liquid scintillation counting at 50% counting efficiency using Ready Protein Plus cocktail (Beckman, Fullerton, CA).

- 5 Nondisplaceable binding was determined in the presence of 1 mM LTB<sub>4</sub> and was usually less than 10% of total binding. Data were analyzed using linear regression analysis of log-logit plots of the values between 10% and 90% of control binding to calculate IC<sub>50</sub>s and slope factors (pseudo-Hill
- 10 coefficients). IC<sub>50</sub> values thus obtained were corrected for radioligand concentration (Cheng and Prusoff, Biochem. Pharmacol., 22, 3099 (1973)) to calculate K<sub>i</sub> values. pK<sub>i</sub> is the mean -log K<sub>i</sub> for n experiments.

- Compounds of the instant invention tested in the above
- 15 assay were found to have a pK<sub>i</sub> of between 7 and 11.

- The ability of compounds of formula I to treat eye inflammation can be evaluated in two models, allergen-induced conjunctivitis in guinea pigs (Garceau et al., European J. Pharmacol., 143, 1-7, 1987) and endotoxin-
- 20 induced uveitis in rats (Okumura et al., Int. Ophthalmol. 14, 31-6, 1990).

## Assay 2

### Allergen-Induced Conjunctivitis Model

- 25 Male guinea pigs, weighing approximately 300 grams are sensitized by injecting intraperitoneally 0.5 ml of a suspension containing 10 µg of ovalbumin and 0.1 gram of aluminum hydroxide in saline. Two weeks later, animals are challenged topically twice, 24 hours apart, on the eye with
- 30 10 µl of a 2.5% (w/v) solution of ovalbumin dissolved in saline. In the absence of an inhibitor, increased microvascular permeability (measured by assaying for the extravasation of a radioactively-labeled protein from the vasculature into the conjunctival tissue) and neutrophil
- 35 infiltration (assayed by measuring myeloperoxidase activity in the conjunctiva) can be observed 30 minutes after the second antigen challenge. Some of the response occurring is

-21-

due to release of histamine and LTD<sub>4</sub>. This can be masked by pretreating the animals 30 minutes before antigen challenge intravenously with mepyramine (2.0 mg/kg) and zafirlukast (1  $\mu$ mole/kg). A compound of formula I in 10 $\mu$ l of vehicle is  
5 applied topically 20 minutes before the second challenge.

### Assay 3

#### Endotoxin-Induced Vreitis Model

In the uveitis model, Salmonella endotoxin (1mg/rat) is  
10 injected into the foot pads of Lewis rats. Eighteen hours later, maximum increases of cells and proteins occur in the anterior chamber in the absence of any anti-inflammatory agent. Three doses of a compound of formula I in 10 $\mu$ l vehicle are applied topically at 5 hour intervals following  
15 injection of the endotoxin. Dose-response effects are obtained by dividing the animals into 4 experimental groups of 10 rats each. The groups are treated with either vehicle, 0.5, 1.0, or 2.0% (w/v) of a compound of formula I in 0.5% hydroxyethyl-cellulose. The effectiveness of a  
20 treatment is accessed by comparing the enhanced fluid and cell infiltration in either the conjunctiva or aqueous humor of the treated group to that of the vehicle control.

Acute iritis can be characterized by moderately severe pain and photophobia with moderately decreased vision.  
25 Transparent precipitates may be present on the posterior surface of the cornea. Intraocular pressure is usually normal or soft and lacrimation is often present along with circumcorneal hyperemia. The iris may appear dull and swollen, the pupil small and irregular with minimal  
30 pupillary response to light.

The therapeutic and prophylactic treatments provided by this invention are practiced by administering to a mammal in need thereof a dose of a compound of formula I or a pharmaceutically acceptable salt or solvate thereof, that is  
35 effective to inhibit or treat iritis.

-22-

The term "inhibit" includes its generally accepted meaning which includes prohibiting, preventing, restraining and slowing, stopping or reversing progression, severity or a resultant symptom. As such, the present method includes  
5 both medical therapeutic and/or prophylactic administration as appropriate.

While it is possible to administer a compound employed in the methods of this invention directly without any formulation, the compounds are usually administered in the  
10 form of pharmaceutical formulation comprising a pharmaceutically acceptable excipient and at least one compound of the present invention. The compounds or formulations of the present invention may be administered by the oral and rectal routes, topically, parenterally, e.g.,  
15 by injection and by continuous or discontinuous intra-arterial infusion, in the form of, for example, tablets, lozenges, sublingual tablets, sachets, cachets, elixirs, gels, suspensions, aerosols, ointments, for example, containing from 0.01 to 90% by weight of the active compound  
20 in a suitable base, soft and hard gelatin capsules, suppositories, injectable solutions and suspensions in physiologically acceptable media, and sterile packaged powders adsorbed onto a support material for making injectable solutions. Such formulations are prepared in a  
25 manner well known in the pharmaceutical art and comprise at least one active compound. See, e.g., REMINGTON'S PHARMACEUTICAL SCIENCES, (16th ed. 1980).

In making the formulations employed in the present invention the active ingredient is usually mixed with an  
30 excipient, diluted by an excipient or enclosed within such a carrier which can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active  
35 ingredient. In preparing a formulation, it may be necessary to mill the active compound to provide the appropriate particle size prior to combining with the other.



-23-

ingredients. If the active compound is substantially insoluble, it ordinarily is milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size is normally adjusted by  
5 milling to provide a substantially uniform distribution in the formulation, e.g. about 40 mesh.

Some examples of suitable carriers, excipients and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate,  
10 alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, tragacanth, gelatin, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil;  
15 wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxybenzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the  
20 active ingredient after administration to the patient by employing procedures known in the art.

The compounds of this invention may be delivered transdermally using known transdermal delivery systems and excipients. Most preferably, a compound of this invention  
25 is admixed with permeation enhancers including, but not limited to, propylene glycol, polyethylene glycol monolaurate, and azacycloalkan-2-ones, and incorporated into a patch or similar delivery system. Additional excipients including gelling agents, emulsifiers, and buffers may be  
30 added to the transdermal formulation as desired.

For topical administration, a compound of this invention ideally can be admixed with any variety of excipients in order to form a viscous liquid or cream-like preparation.

35 For oral administration, a compound of this invention ideally can be admixed with carriers and diluents and molded into tablets or enclosed in gelatin capsules.

-24-

In the case of tablets, a lubricant may be incorporated to prevent sticking and binding of the powdered ingredients in the dies and on the punch of the tableting machine. For such purpose there may be employed for instance aluminum, magnesium or calcium stearates, talc or mineral oil.

Preferred pharmaceutical forms of the present invention include capsules, tablets, creams and ointments. Especially preferred are creams and ointments.

The therapeutic and prophylactic treatments provided by this invention are practiced by administering to a mammal in need thereof a dose of a compound of formula I or a pharmaceutically acceptable salt or solvate thereof that is effective to inhibit or treat iritis.

Advantageously for this purpose, formulations may be provided in unit dosage form, preferably each dosage unit containing from about 5 to about 500 mg (from about 5 to 50 mg in the case of parenteral or inhalation administration, and from about 25 to 500 mg in the case of oral or rectal administration) of a compound of Formula I. Dosages from about 0.5 to about 300 mg/kg per day, preferably 0.5 to 20 mg/kg, of active ingredient may be administered although it will, of course, readily be understood that the amount of the compound or compounds of Formula I actually to be administered will be determined by a physician, in the light of all the relevant circumstances including the condition to be treated, the choice of compound to be administered and the choice of route of administration and therefore the above preferred dosage range is not intended to limit the scope of the present invention in any way.

The specific dose of a compound administered according to this invention to obtain therapeutic or prophylactic effects will, of course, be determined by the particular circumstances surrounding the case, including, for example, the route of administration the age, weight and response of the individual patient, the condition being treated and the severity of the patient's symptoms.

-25-

In general, the compounds of the invention are most desirably administered at a concentration that will generally afford effective results without causing any serious side effects and can be administered either as a  
5 single unit dose, or if desired, the dosage may be divided into convenient subunits administered at suitable times throughout the day.

While all of the compounds illustrated above exemplify LTB<sub>4</sub> inhibition activity in vitro, we have also discovered  
10 that compounds bearing a single acidic group (R<sub>6</sub>) are considerably more orally bioactive when administered to mammals compared with those compounds bearing two such acidic groups. Thus, a preferred embodiment when administering compounds of Formula I orally to mammals  
15 comprises administering compounds bearing a single acidic R<sub>6</sub> functionality.

The following formulation examples may employ as active compounds any of the compounds of this invention. The examples are illustrative only and are not intended to limit  
20 the scope of the invention in any way.

#### Formulation 1

Hard gelatin capsules are prepared using the following  
25 ingredients:

	<u>Quantity (mg/capsule)</u>
30 3-(2-(3-(2-Ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)-6-(4-carboxyphenoxy)phenyl)propanoic acid	250
Starch	200
Magnesium stearate	10

35 The above ingredients are mixed and filled into hard gelatin capsules in 460 mg quantities.

-26-

Formulation 2

A tablet is prepared using the ingredients below:

		<u>Quantity (mg/tablet)</u>
5	1-(4-(Carboxymethoxy)phenyl)-1-(1H-tetrazol-5-yl)-6-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)hexane	250
10	Cellulose, microcrystalline	400
	Silicon dioxide, fumed	10
	Magnesium stearate	5

The components are blended and compressed to form  
15 tablets each weighing 665 mg.

Formulation 3

An aerosol solution is prepared containing the  
20 following components:

		<u>Weight %</u>
25	3-[4-[7-Carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]-9H-xanthene]]propanoic acid	0.25
	Ethanol	30.00
	Propellant 11 (trichlorofluoromethane)	10.25
30	Propellant 12 (Dichlorodifluoromethane)	29.75
	Propellant 114 (Dichlorotetrafluoroethane)	29.75

35 The active compound is dissolved in the ethanol and the solution is added to the propellant 11, cooled to -30°C. and transferred to a filling device. The required amount is then

-27-

fed to a container and further filled with the pre-mixed propellants 12 and 114 by means of the cold-filled method or pressure-filled method. The valve units are then fitted to the container.

5

Formulation 4

Tablets each containing 60 mg of active ingredient are made up as follows:

	2-[2-Propyl-3-[3-[2-ethyl-5-hydroxy-4-(4-fluorophenyl)phenoxy]propoxy]phenoxy]-	
15	benzoic acid sodium salt	60 mg
	Starch	45 mg
	Microcrystalline cellulose	35 mg
	Polyvinylpyrrolidone	4 mg
	(as 10% solution in water)	
20	Sodium carboxymethyl starch	4.5 mg
	Magnesium stearate	0.5 mg
	Talc	1 mg
	Total	150 mg

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders which are then passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50-60° and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

35

-28-

Formulation 5

Capsules each containing 80 mg of medicament are made as follows:

5	5-[3-[2-(1-Carboxy)ethyl]-4-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]-phenyl]-4-pentynoic acid	80 mg
	Starch	59 mg
10	Microcrystalline cellulose	59 mg
	Magnesium stearate	2 mg
	Total	200 mg

The active ingredient, cellulose, starch and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

Formulation 6

20       Suppositories each containing 225 mg of active ingredient are made as follows:

25	3-(2-(3-(2-Ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)-6-(4-carboxyphenoxy)phenyl)propanoic acid	250
	Starch	200
	Magnesium stearate	10

30       The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

-29-

Formulation 7

Suspensions each containing 50 mg of medicament per 5 mL dose are made as follows:

5	2-[2-Propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid	50 mg
	Sodium carboxymethyl cellulose	50 mg
10	Sugar	1 g
	Methyl paraben	0.05 mg
	Propyl paraben	0.03 mg
	Flavor	q.v.
	Color	q.v.
15	Purified water to	5 mL

The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethylcellulose, sugar, and a portion of the water to form a suspension. The parabens, flavor and color are dissolved and diluted with some of the water and added, with stirring. Sufficient water is then added to produce the required volume.

Formulation 8

25 An intravenous formulation may be prepared as follows:


2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid	100 mg
Isotonic saline	1,000 ml

30 The solution of the above ingredients generally is administered intravenously to a subject at a rate of 1 ml per minute.

-30-

We claim:

1. A method for treating or inhibiting iritis in  
 5 a mammal which comprises administering to a mammal in need  
 thereof an effective amount of a compound of the formula I



10

wherein:

15 R<sub>1</sub> is C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>2</sub>-C<sub>5</sub> alkynyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, (C<sub>1</sub>-C<sub>4</sub> alkyl)thio, halo, or R<sub>2</sub>-substituted phenyl;


20 each R<sub>2</sub> and R<sub>3</sub> are each independently hydrogen, halo, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(O)<sub>q</sub>-, trifluoromethyl, or di-(C<sub>1</sub>-C<sub>3</sub> alkyl)amino;

X is -O-, -S-, -C(=O), or -CH<sub>2</sub>-;

Y is -O- or -CH<sub>2</sub>-;

25

or when taken together, -X-Y- is -CH=CH- or



30




Z is a straight or branched chain C<sub>1</sub>-C<sub>10</sub> alkylidenyl;






-31-

5 A is a bond, -O-, -S-, -CH=CH-, or -CR<sub>a</sub>R<sub>b</sub>-, where R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, C<sub>1</sub>-C<sub>5</sub> alkyl, or R<sub>7</sub>-substituted phenyl, or when taken together with the carbon atom to which they are attached form a C<sub>4</sub>-C<sub>8</sub> cycloalkyl ring;

R<sub>4</sub> is R<sub>6</sub> ,



-32-



where,

5 each R<sub>6</sub> is independently -COOH, 5-tetrazolyl, -CON(R<sub>9</sub>)<sub>2</sub>, or -CONHSO<sub>2</sub>R<sub>10</sub>;

each R<sub>7</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>2</sub>-C<sub>5</sub> alkynyl, benzyl, methoxy, -W-R<sub>6</sub>, -T-G-R<sub>6</sub>,  
 10 (C<sub>1</sub>-C<sub>4</sub> alkyl)-T-(C<sub>1</sub>-C<sub>4</sub> alkylidenyl)-O-, or hydroxy;

R<sub>8</sub> is hydrogen or halo;

-33-

each R<sub>9</sub> is independently hydrogen, phenyl, or C<sub>1</sub>-C<sub>4</sub> alkyl, or when taken together with the nitrogen atom form a morpholino, piperidino, piperazino, or pyrrolidino group;

5

R<sub>10</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl or phenyl;

R<sub>11</sub> is R<sub>2</sub>, -W-R<sub>6</sub>, or -T-G-R<sub>6</sub>;

10

each W is a bond or straight or branched chain divalent hydrocarbyl radical of one to eight carbon atoms;

15

each G is a straight or branched chain divalent hydrocarbyl radical of one to eight carbon atoms;

each T is a bond, -CH<sub>2</sub>-, -O-, -NH-, -NHCO-, -C(=O)-, or -S(O)<sub>q</sub>-;

20

K is -C(=O)- or -CH(OH)-;

each q is independently 0, 1, or 2;

p is 0 or 1; and

25

t is 0 or 1;

provided when X is -O- or -S-, Y is not -O-;

30

provided when A is -O- or -S-, R<sub>4</sub> is not R<sub>6</sub>;

provided when A is -O- or -S- and Z is a bond, Y is not -O-; and

35


provided W is not a bond when p is 0;

-34-

or a pharmaceutically acceptable salt or solvate thereof.

2. The method as claimed in **Claim 1** employing a compound of the formula

5



or a pharmaceutically acceptable salt or solvate thereof.

10 3. The method as claimed in **Claim 2** employing 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid or a pharmaceutically acceptable salt or solvate thereof.

15 4. The method as claimed in **Claim 2** employing 3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)-6-(4-carboxy-phenoxy)phenyl)propionic acid or a pharmaceutically acceptable salt or solvate thereof.

20 5. The method as claimed in **Claim 2** employing 1-(4-(carboxy-methoxy)phenyl)-1-(1H-tetrazol-5-yl)-6-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)hexane or a pharmaceutically acceptable salt or solvate thereof.

25 6. The method as claimed in **Claim 2** employing 3-[4-[7-carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]-propoxy]-9H-xanthene]]propanoic acid or a pharmaceutically acceptable salt or solvate thereof.


30 7. The method as claimed in **Claim 2** employing 5-[3-[2-(1-carboxy)-ethyl]-4-[3-[2-ethyl-4-(4-fluorophenyl)-5-

-35-

hydroxyphenoxy]-propoxy]phenyl]-4-pentynoic acid or a pharmaceutically acceptable salt or solvate thereof.

8. The method as claimed in any one of **Claims 1**  
5 to 7 in which the mammal is a human.

9. Use of a compound of the formula I



wherein:


15 R<sub>1</sub> is C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>2</sub>-C<sub>5</sub> alkynyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, (C<sub>1</sub>-C<sub>4</sub> alkyl)thio, halo, or R<sub>2</sub>-substituted phenyl;

20 each R<sub>2</sub> and R<sub>3</sub> are each independently hydrogen, halo, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(O)<sub>q</sub>-, trifluoromethyl, or di-(C<sub>1</sub>-C<sub>3</sub> alkyl)amino;

X is -O-, -S-, -C(=O), or -CH<sub>2</sub>-;

25 Y is -O- or -CH<sub>2</sub>-;

or when taken together, -X-Y- is -CH=CH- or



30


-36-

Z is a straight or branched chain C<sub>1</sub>-C<sub>10</sub> alkylidenyl;



5

A is a bond, -O-, -S-, -CH=CH-, or -CR<sub>a</sub>R<sub>b</sub>-, where R<sub>a</sub> and R<sub>b</sub> are each independently hydrogen, C<sub>1</sub>-C<sub>5</sub> alkyl, or R<sub>7</sub>-substituted phenyl, or when taken together with the carbon atom to which they are attached form a C<sub>4</sub>-C<sub>8</sub> cycloalkyl ring;




R<sub>4</sub> is R<sub>6</sub> ,



10



-37-



where,

5            each R<sub>6</sub> is independently -COOH, 5-tetrazolyl, -CON(R<sub>9</sub>)<sub>2</sub>, or -CONHSO<sub>2</sub>R<sub>10</sub>;

10           each R<sub>7</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>2</sub>-C<sub>5</sub> alkynyl, benzyl, methoxy, -W-R<sub>6</sub>, -T-G-R<sub>6</sub>, (C<sub>1</sub>-C<sub>4</sub> alkyl)-T-(C<sub>1</sub>-C<sub>4</sub> alkylidenyl)-O-, or hydroxy;

R<sub>8</sub> is hydrogen or halo;

-38-

each R<sub>9</sub> is independently hydrogen, phenyl, or C<sub>1</sub>-C<sub>4</sub> alkyl, or when taken together with the nitrogen atom form a morpholino, piperidino, piperazino, or pyrrolidino group;

5

R<sub>10</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl or phenyl;

R<sub>11</sub> is R<sub>2</sub>, -W-R<sub>6</sub>, or -T-G-R<sub>6</sub>;

10

each W is a bond or straight or branched chain divalent hydrocarbyl radical of one to eight carbon atoms;

15

each G is a straight or branched chain divalent hydrocarbyl radical of one to eight carbon atoms;

each T is a bond, -CH<sub>2</sub>-, -O-, -NH-, -NHCO-, -C(=O)-, or -S(O)<sub>q</sub>-;

20

K is -C(=O)- or -CH(OH)-;

each q is independently 0, 1, or 2;

p is 0 or 1; and

25

t is 0 or 1;

provided when X is -O- or -S-, Y is not -O-;

30

provided when A is -O- or -S-, R<sub>4</sub> is not R<sub>6</sub>;

provided when A is -O- or -S- and Z is a bond, Y is not -O-; and

35

provided W is not a bond when p is 0;




-39-

or a pharmaceutically acceptable salt or solvate thereof, optionally in combination with a pharmaceutically acceptable excipient, for the preparation of a pharmaceutical composition for treating or inhibiting iritis in a mammal.

5

10. The use according to **claim 9** employing a compound of the formula



10

or a pharmaceutically acceptable salt or solvate thereof.

11. The use according to **claim 9** wherein the compound employed is 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid

15 or a pharmaceutically acceptable salt or solvate thereof.

12. The use according to **claim 9** wherein the compound employed is 3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)-6-(4-carboxyphenoxy)phenyl)propionic acid or a pharmaceutically

20 acceptable salt or solvate thereof.

13. The use according to **claim 9** wherein the compound employed is 1-(4-(carboxy-methoxy)phenyl)-1-(1H-tetrazol-5-yl)-6-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)hexane or a pharmaceutically acceptable salt

25 or solvate thereof.

-40-

14. The use according to **claim 9** wherein the compound employed is 3-[4-[7-carboxy-9-oxo-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]-propoxy]-9H-xanthene]]propanoic acid or a pharmaceutically acceptable  
5 salt or solvate thereof.

15. The use according to **claim 9** wherein the compound employed is 5-[3-[2-(1-carboxy)-ethyl]-4-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]-propoxy]phenyl]-  
10 4-pentynoic acid or a pharmaceutically acceptable salt or solvate thereof.

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/US98/05414

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/35, 31/28, 31/05, 31/34, 31/44, 31/40, 31/195

US CL : 514/457, 432, 731, 461, 299, 411, 563

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/457, 432, 731, 461, 299, 411, 563

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN: compounds and methods of use

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,462,954 A (BAKER ET AL) 31 October 1995, See columns 1-2, 138 and claims 34-38	1-15

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents	* T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be of particular relevance	* X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* A* document member of the same patent family
* C* document referring to an oral disclosure, use, exhibition or other means	
* P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

06 MAY 1998

Date of mailing of the international search report

23 JUL 1998

Name and mailing address of the ISA-US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

RUSSELL TRAVERS